

# Adult Flat Foot

STANMORE FOOT AND ANKLE SPECIALISTS  
ARE A MULTI-DISCIPLINARY TEAM.

ALL TEAM MEMBERS ARE SPECIALISED  
IN FOOT AND ANKLE CARE AND WORK  
TOGETHER TO PROVIDE AND DELIVER A  
QUALITY SERVICE.

Stanmore Foot  
& Ankle Specialists

part of **HCA** Healthcare UK

# ADULT FLAT FOOT

THE FOLLOWING INFORMATION SHEET HAS BEEN PRODUCED TO PROVIDE OUR PATIENTS WITH AN OVERVIEW OF THEIR CONDITION AND WHAT TREATMENTS THEY MAY EXPECT. HOWEVER, EACH PERSON IS DIFFERENT AND THE TREATMENT YOU ARE OFFERED WILL DEPEND ON YOUR UNIQUE CIRCUMSTANCES.

## What is a flat foot deformity?

Most feet have an arch on the inside of the foot. In some people this arch is less obvious and the foot looks 'flatter'. Up to 1 in 5 people may have a slightly flat foot. In some people this is the normal shape of their foot, but in others the deformity can develop over time. A flat foot is not always a problem but if you are getting pain or your foot gets flatter over time, it may need treatment.

## What causes a flat foot deformity?

In some people this is just the normal shape of your foot and may not cause any problems. Some people have a flat foot which is caused due to imbalances due to a nerve or muscle problems or due to a developmental problem. However, in most people with an acquired deformity some of the soft tissue structures on the inside of the foot begin to wear out.

This may involve a tendon called the tibialis posterior tendon and a structure called the spring ligament which can wear out with time. These structures normally support the arch of the foot and when damaged they can get swollen and painful or may even tear in severe cases. When this happens the shape of the foot can get worse.

There are other factors which can play a role in developing a flat foot including injury to the tendon, increasing age and increasing weight.





## What Symptoms does a flat foot cause?

Most people with flat feet will have very few problems. When problems occur, it usually starts with a pain on the inside of the foot. This may start as a dull ache on walking longer distances but may become worse over time.

### Other common symptoms include:

#### Pain

Pain usually starts on the inside of the foot, but may be felt along the back of the ankle as well along the path of the tendon. If the tendon becomes inflamed then this may be associated with swelling and a burning or shooting pain. If the deformity is more severe then pain may be felt on the outside of the ankle or in the ankle itself.

#### Difficulty walking

It may become difficult to walk long or even short distances and pain may become worse at the end of the day.

#### Change in foot shape

As the foot becomes flatter it may also start to point outwards more. You may notice that your shoes begin to wear out more easily because of this.

#### Inability to stand on tip toes

As the tendon gets weaker and more painful you may find that you have difficulty standing on tip toes.





# Non-Operative Treatment

## Treatment for Adult Flat Foot

Most people with flat foot do not need surgery. The exact treatment will depend on your symptoms and the severity of your deformity. If you are having symptoms common things which can be tried are:

### RICE - Rest, Ice, Elevation, Compression

Resting your foot and reducing the amount you walk can help for acute episodes of pain. This includes refraining from sports and impact activities which you are healing. You can also apply ice to help reduce the inflammation. It is important not to apply ice directly as this can damage your skin. A tubigrip or elasticated support can provide some support and pain relief. Elevating the foot can also help to reduce swelling.

### Pain Killers

In the acute setting, pain killers such as non-steroidal anti-inflammatories can help. These can be applied as a gel or cream on the foot or taken as tablets.

### Brace or boot

If you are in a lot of pain you may benefit from a short period of time in a special boot or brace for a up to 6 weeks.

### Insoles

In the longer term, specially designed insoles can help to support the arch of your foot and take the pressure off your tendon. This can help improve your symptoms and may help stop your deformity from getting worse.

### Shoes

It is also important to wear flat shoes which support your foot. Lace-up shoes may help to secure your foot and keep the insoles in place.

### Physiotherapy

Physiotherapy is aimed at stretching your achilles tendon, and improving the strength and balance of the muscles in the foot and ankle. It can also help to balance the load in your tibialis posterior tendon which helps it heal.







# Surgery for Flatfoot

## When might you need surgery?

If the above measures haven't helped, the surgery may be an option. Also if you have a progressive deformity or begin to get arthritis then surgery may be advised.

## Benefits of surgery

Surgery aims to:

- Treat the pain associated with a flat foot deformity
- Correct the deformity
- Stop the deformity from progressing


In order to feel the full benefits from surgery it may take up to a year.

## What does the procedure involve?

There are many different procedures which may be done for treating flat feet. These range from:

- Debridement of the tendon which is inflamed
- Correction of the deformity by realigning your foot and reconstructing the spring ligament and tibialis posterior tendon
- Fusing bones in your foot together to treat the deformity and arthritis

The exact procedure will be discussed with you by your surgeon, but in general one or more cuts are made around your foot and ankle. The bones and joints are shifted to balance the foot. The bones are put together and held in place with either screws or screws and a plate.



You may also need to have a tendon which bends your toes taken to help support the arch of your foot. Other devices and implants may also be used to support the operation. After the operation the skin is closed, and you will be placed into a cast. You will not be allowed to put weight through your leg for about 4 to 6 weeks. You may be in a boot for a further 6 weeks after this.

## What are the risks of the procedure?

- All operations have risks
- Any medical conditions you already have may become worse due to the operation.
- Some risks may be rarer but more relevant to you. It is important to discuss with your surgeon your particular circumstances.

## Common Risks (Occur in up to 5 out of 100 procedures)

### Pain

For most operations, you will be given a local anaesthetic block to reduce pain, but you may still have some pain or discomfort. Usually, this can be controlled with simple pain management tablets.

### Swelling

Feet tend to swell up because of gravity, and this can last for some months.

### Scarring

Any type of surgery will leave a scar, sometime this can be painful or sensitive.

### Minor wound redness

With all operations there is a risk of infection. Sometimes the wound edges can become a bit red or may not fully heal. In some cases you may require antibiotics to get this to settle. Risks are higher if you have diabetes, if you smoke, or if you are on medication which alters your immune system (eg steroids or rheumatoid medication).

### Numbness

After surgery you may have some numbness or tingling around the scar. This is because small nerves to the skin may have been cut. This usually recovers with time, but sometimes may be permanent.



### Prominent metalwork

In some cases the screws or plates can be felt under the skin, requiring a second procedure to remove them once the bones are healed.

### Problematic weight distribution

Because foot and ankle surgery shifts the balance of weight across the foot, pain can appear in areas of the foot that previously did not have pain. Most cases settle with physiotherapy and management with insoles, but further surgery might be necessary.

## Less Common Risks (occur in up to 1 in 100 procedures)

### Blood clots

Because you will not be allowed to walk immediately after the surgery blood clots can occur and can lead to swelling of the leg (deep vein thrombosis) or chest pain (pulmonary embolism).

### Non-union

This is when the bones do not heal to one another. Factors that have been shown to prevent bone healing include smoking, vitamin D deficiency, thyroid disease and diabetes.

### Malunion

This is when the bones do not heal in the desired position. Sometimes this can cause discomfort and rubbing on shoes. Rarely, another procedure may be required to correct this.

### **Implant loosening**

Any implants used can loosen on occasion. Although the cause can be unknown, it can be related to infection, rejection of the implant or failure of bones to join.

### **Overcorrection**

The correction involves balancing the bones and soft tissues. Sometimes the deformity can be overcorrected. This complication often requires further surgery.

### **Recurrence of deformity**

Correcting a deformity involves balancing the foot and ankle. Sometimes, the balancing fails. If this happens, further surgery may be required.

## **Rare risks**

**(occur in less than 1 in a 100 procedures)**

### **Deep infection**

Although every precaution is taken to prevent infection, it can still happen. If the infection does not settle with antibiotics, you may need a further operation to remove any metal and clear the infection.

### **Intraoperative fracture or broken metalwork**

rarely a fracture may occur during surgery or a metal pin or screw could break. The surgeon will act in your best interests at the time of surgery to give you the best outcome.

### **Complex regional pain syndrome**

This is where the "fight or flight" nerves that supply the foot become over active and this can cause swelling, stiffness, pain, and colour and temperature changes to the foot. Treatment involves physiotherapy and other treatments and it could take several months to improve.

### **Nerve injury**

If a larger nerve supplying the toe / foot becomes damaged or caught in scar tissue, it could lead to ongoing pain, numbness and tingling. Such damage is usually temporary and the sensation returns over some time, but it can be permanent.

### **Tendon injury**

During the operation, a tendon may be injured or can become caught in the scar leading to reduced movement. Rarely this requires further surgery.

### **Blood vessel damage**

If the blood supply to a toe is damaged it could lead to part or all of the toe becoming permanently damaged. In severe cases, the damaged parts of the toe may need to be removed.

### **Rupture**

On occasion, the tendon procedure can result in the tendon being weak and rupturing. This may require treatment in a cast/boot or even surgery.

### **Compartment Syndrome**

This is a rare condition where the muscles in either the lower leg or foot swell. If this did occur it is very painful and would likely require emergency treatment. The most common cause is a tight cast which would need to be removed, however if this did not resolve the problem surgery would be required.

### **Amputation**

Although extremely rare, loss of limb can result from surgery especially if deep infection or blood vessel injury occurs. The risk for this complication is increased in diabetics and smokers.

### **Death**

Whilst this is extremely rare for foot and ankle surgery, it can occur especially if there are pre-existing medical conditions.



# Before & After Surgery

## What happens before the procedure?

Before the operation you will need to have a pre-admission assessment to ensure that you are medically fit for surgery. You also need to consider a number of other practical points to prepare yourself for surgery:

- Can someone help you carry out basic every day tasks such as preparing food and food shopping?
- If you have stairs, how will you get up and down them? Do you have sturdy hand rails? If your toilet is downstairs, would it be easier to have your bed downstairs until you have recovered and are able to negotiate the stairs safely?
- Will you be able to manage crutches, or will you need to hire a wheelchair or a knee scooter?
- If you plan on taking a shower you will need to be able to get in and out of a bath safely and will need to keep your cast dry – there are a number of covers which are specially designed for casts that you can buy in shops or online.
- Do you have special pain management needs?

Some of these will be discussed with you in pre-assessment and you may also need to be assessed by a physiotherapist or occupational therapist if you have any concerns about coping at home after the procedure. The better prepared you are the smoother the whole process will be and the less time you will need to spend in hospital.

## On the day of surgery

Surgery for flat foot usually requires at least an overnight stay and you may need to stay up to 2 nights. You will need to bring with you a bag with essential toiletries, phone / table / laptop chargers, and other clothing items you may need for your stay.

You will also need to have a sturdy shoe for your other foot to make sure you can mobilise normally. If you have orthotics or braces for your other side, or other walking aids, please make sure you bring these with you.

On the day of surgery you will be first be admitted onto the ward. You will see a number of members from the nursing team, and also the surgical and anaesthetic teams. You will have been told when you could last eat and drink and it is important that you do not eat or drink after the times that have been told to you.



You will have an arrow/mark put on the leg by your surgeon and you will sign a consent form confirming you agree to surgery. You will also have the opportunity ask any further questions to the surgeon or anaesthetist.

When it is time for your surgery you will be taken down to theatre and there you will be given an anaesthetic (the type of anaesthetic will be discussed with you beforehand). Most flat foot surgery is done with a nerve block in addition to the anaesthetic to help with post-operative pain control. After surgery you will be taken to a recovery room and when you are fully awake you will go back to your ward.

On the ward you will be given pain killers and other medication you require, including injections or tablets to help prevent blood clots. The next day you will be seen by physiotherapists to help you mobilise and you will be reviewed by a member of the surgical team. Once you are fit to go home any pain killers, blood clot preventers and other medication will be sorted out for you and you will be discharged if it is safe to do so.

## After your surgery

You will have a number of appointments to be made for you after your surgery. The following are rough guidelines, but every case will be different and you may be given specific instructions according to the exact procedure you have had. If you have had fusions then you may be in cast for longer.

For the first 2 weeks it is very important that you keep walking around to a minimum and keep your leg elevated. It needs to ideally be above or at the level of your heart (or as close as possible) for 23 hours a day. This means that every hour you can get up for about 5 minutes to get food/drink, or go to the toilet/shower, etc. This may mean that you need to keep your leg elevated on pillows in bed or on a stool when sitting. Keeping it elevated is important to reduced swelling, reduce pain and let the wound heal well. You are encouraged to move your toes to keep the circulation going and you will be given blood thinning injections or tablets to use for 6 weeks.



You will then usually be seen at around 2 weeks. The temporary cast will be removed and your wounds inspected. If appropriate the stitches will be removed /trimmed and you are placed into another, full cast. At this stage you should still keep your weight off your foot, but you can usually move around a bit more.

At 6 weeks after your surgery you are seen in clinic and the cast is removed and you will have x-rays. If all looks well you can usually go into a removable boot which you may need for the next 4 to 6 weeks. You may also begin some gentle physiotherapy at this stage depending on your progress.

At 3 months if all is well you can begin to mobilise freely and go into your normal shoes. At this stage you will likely need a new insole to support the foot that has been repaired. It does however take about 6 months to a year for you to recover fully from this operation.

## Things to look out for

When at home there are a few things you should look out for. You should contact the surgical team if you have any of the following problems:

- Increasing swelling
- Cast which starts to feel tight
- Pain which is getting worse
- Pins and needles

If you are unable to contact the surgical team, or for example it is out of hours and your symptoms are getting worse you should go to your local emergency department. You should also go to your emergency department or call an ambulance if you have any of the following symptoms as they may indicate potentially serious problems:

- Chest pain
- Sudden shortness of breath
- Severe progressive pain despite elevation
- Severe swelling of your leg/thigh
- You become hot, clammy and dizzy

## Getting back to normal

Once again, these are rough guidelines and your particular situation may differ. In general:

### Work

If your job is mostly sitting you may be able to return to work 4 weeks following your surgery providing you can get to work safely and you can keep your leg elevated. If your job involves walking or standing, then this may take longer, perhaps up to 8 weeks, or even longer if you have had a fusion

### Walking

Most people should be able to walk without aids about 3 months following surgery, but it may take some time to build up your walking distance and this will depend on how well you walked before the surgery

### Shoes

It can take a number of months for the swelling to fully settle after surgery. It is normal to have swelling by the end of the day for up to 6 months or sometimes more following surgery. By 6 months most patients can go back into most of their normal footwear

### Driving

If you have an automatic car and have had left foot surgery, you can normally drive 2 weeks after your surgery, providing you are driving short distances and you can get in and out of the car safely. Otherwise it may take at least 3 months to be able to drive safely. In order to be able to drive you need to be able to safely perform an emergency stop. It is your responsibility to inform your insurance company about the operation you have had to ensure your insurance remains valid.

### Sport

This will depend on the exact operation you have had and what sport you are trying to participate in. Generally you can get back to low impact sports by about 6 months following your surgery.





## Our group

The Harley Street Clinic  
The Lister Hospital  
London Bridge Hospital  
The Portland Hospital  
The Princess Grace Hospital  
The Wellington Hospital  
The Christie Private Care  
HCA UK at University College Hospital  
HCA UK at The Wilmslow Hospital  
Leaders in Oncology Care

To book an appointment with the Stanmore Foot and Ankle Specialists please contact:

**Tel:** +44(0)207 374 7327

**Email:** [sfas@hcahealthcare.co.uk](mailto:sfas@hcahealthcare.co.uk)

**Stanmore Foot  
& Ankle Specialists**

---

part of **HCA**Healthcare UK